Correction

Correction: A novel derivative of betulinic acid, SYK023, suppresses lung cancer growth and malignancy

Tsung-I Hsu^{1,2}, Ying-Jung Chen², Chia-Yang Hung³, Yi-Chang Wang³, Sin-Jin Lin², Wu-Chou Su⁴, Ming-Derg Lai^{1,3,5}, Sang-Yong Kim⁶, Qiang Wang⁶, Keduo Qian⁶, Masuo Goto⁶, Yu Zhao⁶, Yoshiki Kashiwada⁷, Kuo-Hsiung Lee^{6,8}, Wen-Chang Chang^{1,2,3,9,10} and Jan-Jong Hung^{1,2,3,9,10}

Published:

Copyright: Hsu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been corrected: The control data used in the assembly of Figure 5 was incorrect. After reviewing all of the raw data, an updated Figure 5 was created and is shown below. The authors declare that these corrections do not change the results or conclusions of this paper.

Original article: Oncotarget. 2015; 6:13671–13687. https://doi.org/10.18632/oncotarget.3701

¹Center for Infection Disease and Signal Research, College of Medicine, National Cheng Kung University, Tainan, Taiwan

²Institute of Bioinformatics and Biosignal Transduction, College of Bioscience and Biotechnology, National Cheng Kung University, Tainan, Taiwan

 $^{^{3}}$ Institute of Basic Medical Sciences, College of Medicine, National Cheng Kung University, Tainan, Taiwan

⁴Department of Internal Medicine, College of Medicine and Hospital, National Cheng Kung University, Tainan, Taiwan

⁵Department of Biochemistry and Molecular Biology, College of Medicine, National Cheng Kung University, Tainan, Taiwan

⁶Natural Products Research Laboratories, UNC Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, NC, USA

⁷Laboratory of Pharmacognosy, Graduate School of Pharmaceutical Sciences, The University of Tokushima, Tokushima, Japan

⁸Chinese Medicine Research and Development Center, China Medical University and Hospital, Taichung, Taiwan

⁹Department of Pharmacology, College of Medicine, National Cheng Kung University, Tainan, Taiwan

¹⁰Graduate Institute of Medical Sciences, College of Medicine, Taipei Medical University, Taipei, Taiwan




Figure 5: Effects of BA and SYK023 on lung tumor metastasis in vitro and in vivo. (A) After BA and SYK023 treatment for 36 h, H1299 cells were subjected to transwell invasion and wound-healing assay. (B) Representative images of lungs from SCID mice with metastasis (a). The number of surface lung tumors. Data are expressed as mean \pm s.e.m, *P*-value is indicated (b). (C) H1299 cells on the coverslip were treated with the indicated drug, and subjected to immunofluorescent staining for F-actin and DAPI (1000×).