

Endothelial integrin $\alpha 3\beta 1$ stabilizes carbohydrate-mediated tumor/endothelial cell adhesion and induces macromolecular signaling complex formation at the endothelial cell membrane

Olga V. Glinskii^{1,2}, Feng Li^{1,2}, Landon S. Wilson³, Stephen Barnes³, Kate Rittenhouse-Olson⁴, Joseph J. Barchi, Jr.⁵, Kenneth J. Pienta⁶, Vladislav V. Glinsky^{1,2}

¹ Research Service, Harry S. Truman Memorial Veterans Hospital, Columbia, MO

² Department of Pathology and Anatomical Sciences and the Department of Medical Pharmacology and Physiology, University of Missouri School of Medicine, Columbia, MO

³ Department of Pharmacology and Toxicology and Targeted Metabolomics and Proteomics Laboratory, University of Alabama at Birmingham, Birmingham, AL

⁴ Department of Biotechnical and Clinical Laboratory Sciences and the Department of Microbiology, University of Buffalo, Buffalo, NY

⁵ Chemical Biology Laboratory, Center for Cancer Research, National Cancer Institute, Frederick National Laboratory for Cancer Research, Frederick, MD

⁶ Department of Urology, The James Buchanan Brady Urological Institute, Departments of Oncology and Pharmacology and Molecular Sciences, The Johns Hopkins School of Medicine, Baltimore, MD

Correspondence to: Vladislav V. Glinsky, **email:** glinskiivl@health.missouri.edu

Keywords: tumor metastasis, adhesion, Thomsen-Friedenreich antigen, galectin, integrin

Received: February 10, 2014

Accepted: March 19, 2014

Published: March 20, 2014

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT:

Blood borne metastatic tumor cell adhesion to endothelial cells constitutes a critical rate-limiting step in hematogenous cancer metastasis. Interactions between cancer associated carbohydrate Thomsen-Friedenreich antigen (TF-Ag) and endothelium-expressed galectin-3 (Gal-3) have been identified as the leading molecular mechanism initiating tumor/endothelial cell adhesion in several types of cancer. However, it is unknown how these rather weak and transient carbohydrate/lectin mediated interactions are stabilized. Here, using Western blot and LC tandem mass spectrometry analyses of pull-downs utilizing TF-Ag loaded gold nanoparticles, we identified Gal-3, endothelial integrin $\alpha 3\beta 1$, Src kinase, as well as 5 additional molecules mapping onto focal adhesion pathway as parts of the macromolecular complexes formed at the endothelial cell membranes downstream of TF-Ag/Gal-3 interactions. In a modified parallel flow chamber assay, inhibiting $\alpha 3\beta 1$ integrin greatly reduced the strength of tumor/endothelial cell interactions without affecting the initial cancer cell adhesion. Further, the macromolecular complex induced by TF-Ag/Gal-3/ $\alpha 3\beta 1$ interactions activates Src kinase, p38, and ERK1/2, pathways in endothelial cells in a time- and $\alpha 3\beta 1$ -dependent manner. We conclude that, following the initial metastatic cell attachment to endothelial cells mediated by TF-Ag/Gal-3 interactions, endothelial integrin $\alpha 3\beta 1$ stabilizes tumor/endothelial cell adhesion and induces the formation of macromolecular signaling complex activating several major signaling pathways in endothelial cells.

INTRODUCTION

Metastasis is a major cause of cancer-related morbidity and mortality. The initial arrest of blood borne

metastatic tumor cells in distant organ microvasculature constitutes an important rate-limiting step in hematogenous cancer metastasis [1-3]. Consequently, there has been a substantial surge in recent years in research aiming to identify the molecules mediating

tumor cell adhesion to endothelial cells [4-13]. Among others, interactions between cancer-associated carbohydrate Thomsen-Friedenreich antigen (TF-Ag), core 1 disaccharide Gal β 1-3GalNAc expressed on 80-90% of all adenocarcinomas [14-16], and endothelium expressed ~30kDa β -galactoside binding lectin galectin-3 (Gal-3) have been identified as the leading molecular mechanism mediating the initial stages of adhesion between metastatic cells and endothelium in multiple types of cancer including breast [9, 17-19], prostate [9, 17, 20], colon [7] and pancreas [6]. The multifaceted role for Gal-3 in tumor metastasis was first established by the pioneering works from the group of Dr. Avraham Raz [21-24]. Subsequently, it has been demonstrated that, while cancer cell expressed Gal-3 is involved in regulation of tumor cell apoptosis and homotypic aggregation [9, 21, 24, 25] and may even serve as a complementary serum marker in prostate cancer [26], it is endothelium expressed Gal-3 that mediates metastatic cell adhesion to the endothelium via interactions with cancer associated TF-Ag [9, 17-20]. Specifically, it has been shown that in the process of metastasis-associated endothelial activation TF-Ag expressed on either circulating tumor-associated glycoproteins or blood borne metastatic cells induces Gal-3 translocation to and clustering at the endothelial cell membrane [17, 27]. Subsequently, TF-Ag expressing cancer cells interact with cell surface clustered endothelial Gal-3 to initiate metastatic cell arrest in distant organ vasculature [9, 17-20]. However, these carbohydrate/lectin mediated adhesive interactions are rather weak and transient in nature. Without further stabilization transiently adhered cancer cells could be dislodged by the shear force of the flowing blood resulting in the interruption of the metastatic cascade. Indeed, Gal-3 does not contain a transmembrane domain and, therefore, lacks means of being anchored at the endothelial cell surface without the involvement of additional molecules capable of providing such anchorage. However, it is currently unknown which molecules stabilize tumor/endothelial cell adhesion downstream of TF-Ag/Gal-3 interactions.

Recently, it has been shown that, similarly to cancer-associated TF-Ag, NG2 proteoglycan produced by pericytes mediates Gal-3 translocation to and clustering at the endothelial cell membrane followed by the mobilization of endothelial α 3 β 1 integrin, which physically interacts with Gal-3 scaffolds [28]. This information led us to hypothesize that endothelial integrins such as α 3 β 1 could be likewise mobilized to Gal-3 clustered at the endothelial cell membrane during TF-Ag mediated metastatic cell adhesion to the endothelium and stabilize initial transient carbohydrate/lectin mediated interactions. In the present study, we report that endothelial integrin α 3 β 1 does indeed interact directly with TF-Ag/Gal-3 complexes and significantly increases the strength of tumor/endothelial cell adhesion. Further, upon tumor cell/endothelial cell interactions, we

detected activation of several major signaling pathways downstream of endothelial α 3 β 1 suggesting that TF-Ag/Gal-3 mediated interactions induce complex signaling crosstalk in endothelial cells, which could be potentially targeted for therapeutic purposes.

RESULTS

Endothelial integrin α 3 β 1 associates physically with TF-Ag/Gal-3 clusters and induces the formation of focal adhesion type macromolecular signaling complexes

Previously, we demonstrated that in the process of metastasis-associated endothelium activation TF-Ag expressed on circulating tumor-associated glycoproteins or blood borne metastatic cells induces Gal-3 translocation to and clustering at the endothelial cell outer membranes [17, 27]. Subsequently, TF-Ag expressing tumor cells interact with cell surface clustered endothelial Gal-3 to initiate metastatic cell adhesion to the vascular wall [9, 17-20, 27]. To identify integrin molecules and other proteins interacting with TF-Ag/Gal-3 complexes at the endothelial cell membrane we have performed a pull-down assay using gold nanoparticles bearing multiple TF-Ag moieties (TF-Au) covalently attached to the particles through the polyethylene glycol (PEG) linker and control nanoparticles, on which PEG linkers are terminated with the OH groups (PEG-Au) as described in Materials and Methods. Consistent with TF-Ag interactions with Gal-3 scaffolds, Western blot analysis of the pull-down isolates confirmed the presence of Gal-3 in TF-Au samples (Fig. 1, A). Probing the membranes with the panel of antibodies directed against α 3, α 4, α 5, α V, β 1, β 3, β 4, and β 5 integrins revealed the presence of α 3 and β 1 integrin subunits (i.e. α 3 β 1 integrin) in TF-Au, but not in untreated endothelial cell or PEG-Au samples (Fig. 1, A) indicative of α 3 β 1 interaction with TF-Ag/Gal-3 complexes. The signals for α 5 and β 4 integrins were clearly identifiable as well (Fig. 1, A). However, they were present in both TF-Au and PEG-Au samples suggesting that they were pulled down due to nonspecific interactions with PEG. In addition, a principal signal transducing molecule downstream of α 3 β 1 integrin, the Src kinase, was also present in TF-Au pull-downs (Fig. 1, A). Next, individual bands were cut from NuPAGE 4-12% gradient Bis-Tris gels stained with modified colloidal Coomassie Blue G-250 (Fig. 1, B) and analyzed by LC-tandem mass spectrometry following in gel digestion and tryptic peptide extraction. Using this approach, we identified with high fidelity (MOWSE scores ranging from 158 to 1843) additional 5 proteins, which were present in TF-Au samples only: filamin B, β (Mr 263,856); talin (Mr 269,486); vinculin isoform VCL (Mr 116,649); zyxin (Mr

61238); and plastin-3/T-plastin (Mr 70,391) in bands 2, 3, 7, 12, and 13 respectively (Fig. 1, B). Remarkably, all of these 5 proteins map onto focal adhesion pathway. Taken together, these results strongly suggest that interactions of tumor-associated TF-Ag with endothelial Gal-3 mobilize endothelial $\alpha\beta 1$ integrin, which associates physically with Gal-3 scaffolds and induces the formation of focal adhesion type macromolecular signaling complexes at the endothelial cell membrane. Thus, our next two questions were: (i) Whether $\alpha\beta 1$ integrin engagement stabilizes tumor/endothelial cell adhesion; and (ii) Whether these interactions induce major signal transduction pathways in endothelial cells downstream of $\alpha\beta 1$ integrin?

Integrin $\alpha\beta 1$ stabilizes tumor/endothelial cell adhesion

Here, we have employed a modified parallel flow chamber assay to analyze temporal dynamics of changes in the strength of adhesion between metastatic tumor cells and endothelial monolayers. The assay has been designed to allow tumor cells to interact with endothelial monolayers for various time periods under static conditions followed by the application of the increasing wall shear force of a defined magnitude in a parallel flow chamber to displace stably adhered tumor cells. In control experiments (Fig. 2, A), we observed a significant time-dependent increase in a wall shear force necessary to displace 50% of stably adhered tumor cells rapidly reaching the values far exceeding physiological wall shear stress (4-8 dynes/cm²) at which tumor cell adhesion to the vascular wall occurs.

In our experiments, after 10 min of incubation the wall shear stress displacing 50% of tumor cells was 56.4 ± 12.0 dynes/cm² and it was reaching upwards of 250 dynes/cm² after 30 min (Fig. 2, A). These results indicate that tumor/endothelial cell adhesion undergoes rapid stabilization resulting in a significant increase in the adhesion forces between the two cell types. To investigate whether $\alpha\beta 1$ integrin is responsible for the observed tumor/endothelial cells adhesion stabilization, we used anti- $\alpha\beta 1$ function blocking antibody P1B5 [29]. Performing experiments in the presence of 10 $\mu\text{g/ml}$ of anti- $\alpha\beta 1$ did not change significantly the number of stably adhered tumor cells compared with control mouse IgG (Fig. 2, B). However, the wall shear stress required to displace 50% of tumor cells has been significantly ($p < 0.05$) reduced by anti- $\alpha\beta 1$ from 56.4 ± 12.0 dynes/cm² to 9.2 ± 4.3 dynes/cm² (Fig. 2, C). Further, the percent of tumor cells displaced after 10 min of incubation by the flow inflicting wall shear stress of 20 and 40 dynes/cm² has been also increased significantly by anti- $\alpha\beta 1$ antibody (Fig. 2, D). Taken together, these results demonstrate that $\alpha\beta 1$ does not affect the initial (carbohydrate-mediated) tumor cell adhesion to endothelial cells, but plays a critical role in stabilizing tumor/endothelial cell adhesive interactions.

Tumor/endothelial cell interactions activate major signaling pathways in endothelial cells in $\alpha\beta 1$ dependent manner

As our TF-Gold pull-down experiments revealed, in addition to Gal-3 and $\alpha\beta 1$ integrin, the presence of




Figure 1: Macromolecular complexes formed at the endothelial cell membrane downstream of TF-Ag/Gal-3 interactions.

A, Western blot analysis of the integrin molecules and other proteins interacting with TF-Au nanoparticles at the endothelial cell surfaces. Note specific interactions of Gal-3 and integrin subunits $\alpha 3$, and $\beta 1$ with TF-Au nanoparticles indicative of $\alpha\beta 1$ interactions with TF-Ag/Gal-3 complexes. The Src kinase, a principal signal transduction molecule downstream of $\alpha\beta 1$ integrin also interacts specifically with TF-Au nanoparticles. B, Using LC-tandem mass spectrometry, additional 5 proteins mapping onto focal adhesion pathway [filamin B, β (Mr 263,856); talin (Mr 269,486); vinculin isoform VCL (Mr 116,649); zyxin (Mr 61238); and plastin-3/T-plastin (Mr 70,391)] were identified in bands 2, 3, 7, 12, and 13 respectively in TF-Au, but not PEG-Au or control endothelial cell (EC) pull-downs. In B, bands corresponding to heavy chain and light chain of the anti-TF-Ag antibody used for a pull-down denoted as HC and LC respectively. The experiments were performed three times for Western analysis and twice for LC-tandem mass spectrometry with the same results.

the Src kinase (a principal signal transduction molecule acting downstream of $\alpha 3\beta 1$), as well as of 7 other proteins mapping onto focal adhesion pathway (Fig. 1) in TF-Au samples, our next question was whether tumor/endothelial cell adhesive interactions activate major signaling pathways in endothelial cells downstream of $\alpha 3\beta 1$ integrin. Typically, $\alpha 3\beta 1$ induces via Src kinase major MAPK (p38 and ERK1/2) signal transduction pathways. Indeed, Western blot analysis of the endothelial cell lysates following tumor/endothelial cell co-culture experiments

demonstrated time-dependent activation of Src, p38 MAPK, and MEK1/2 in endothelial cells, when they interact with tumor cells (Fig. 3, A). Performing the same experiments in the presence of the function-blocking anti- $\alpha 3\beta 1$ antibody P1B5 abolished time-dependent activation of these signaling pathways (Fig. 3, B). These results indicate that tumor/endothelial cell interactions induce activation of Src and major MAPK signal transduction pathways (p38 and ERK1/2) in endothelial cells in a time-dependent manner downstream of endothelial $\alpha 3\beta 1$




Figure 2: Endothelial integrin $\alpha 3\beta 1$ stabilizes tumor/endothelial cell adhesion. A, Tumor/endothelial cell adhesion undergoes rapid stabilization. Note rapid increase in a wall shear stress required to displace 50% of stably adhered tumor cell after 10, 15, 20, and 30 min of incubation with endothelial monolayers. B, Function blocking anti- $\alpha 3\beta 1$ antibody P1B5 (blue bar) does not affect the initial adhesion of tumor cells to the endothelium after 10 min of incubation compared with control IgG (black bar). C and D, Anti- $\alpha 3\beta 1$ function blocking antibody P1B5 inhibits tumor/endothelial cell adhesion stabilization. C, Compared with control IgG (black bar), anti- $\alpha 3\beta 1$ antibody P1B5 (blue bar) reduces significantly the wall shear stress required to displace 50% of stably adhered tumor cell after 10 min of incubation. D, Compared with control IgG (black bars), anti- $\alpha 3\beta 1$ antibody P1B5 (blue bars) increases significantly the percent of tumor cells displaced by the wall shear stress of 20 and 40 dynes/cm² after 10 min of incubation. In A through D, data presented as means \pm STDEV; *denotes statistical significance ($p < 0.05$)




Figure 3: Activation of major signaling pathways in endothelial cells downstream of integrin $\alpha 3\beta 1$. A, Time-dependent activation of Src, p38, and MEK1/2 in endothelial cells following their interaction with tumor cells for the indicated time periods. B, Anti- $\alpha 3\beta 1$ function blocking antibody P1B5 inhibits time-dependent activation of Src, p38, and MEK1/2 in endothelial cells following their interaction with tumor cells for the indicated time periods. The experiments were repeated at least three times for each setting with similar results.

integrin.

DISCUSSION

The results presented in this study add important new insights to our understanding of the extremely complex multi-step process of hematogenous cancer metastasis. They demonstrate that, following the initial transient adhesive interactions between metastatic tumor cells and vascular endothelium mediated by cancer-associated TF-Ag and endothelium expressed Gal-3, endothelial integrin $\alpha 3\beta 1$ physically associates with TF-Ag/Gal-3 complexes at the endothelial cell surfaces and stabilizes tumor/endothelial cell adhesion. This stabilization results in a rapid increase of the strength of adhesion between endothelial and tumor cells reaching the values far exceeding physiological shear forces acting upon adhered metastatic cells in the circulation. Without such stabilization, metastatic tumor cells attached to the vascular wall via weak carbohydrate/lectin interactions will be dislodged by the flowing blood and metastatic cascade will be interrupted. This makes the endothelial $\alpha 3\beta 1$ integrin an attractive therapeutic target for controlling metastatic spread of cancer.

Further, in addition to stabilizing tumor/endothelial cell adhesive interactions, endothelial $\alpha 3\beta 1$ engagement triggers the formation of the focal adhesion type macromolecular signaling complexes at the endothelial cell membrane and time-dependent phosphorylation of Src, p38, and MEK1/2 in endothelial cells downstream of $\alpha 3\beta 1$ indicative of the activation of p38 and ERK1/2 signal transduction pathways. As these pathways regulating important cellular functions related to cell survival, motility, and proliferation share multiple downstream targets, these results show that tumor/endothelial cell interactions induce complex crosstalk between major MAPK signal transduction pathways, which could have a profound effect on endothelial cell behavior. The biological outcomes of this crosstalk in a context of cancer metastasis and its potential effects on subsequent steps of the metastatic cascade such as tumor cell transendothelial migration and extravasation are currently under investigation and could yield yet additional therapeutic targets for anti-metastatic interventions. For example, previous studies from the group of Dr. M. Sharon Stack have shown that, when both p38 and ERK1/2 MAPK signaling pathways are simultaneously activated in tumor cells downstream of $\alpha 3\beta 1$ integrin, the resulting outcome of the interplay between the two pathways is the activation of tumor cell MMP system in a p38 dependent manner [30]. Whether the same scenario takes place when p38 and ERK1/2 pathways are activated downstream of $\alpha 3\beta 1$ in endothelial cells as a consequence of tumor/endothelial cell adhesion, and whether endothelial p38, ERK1/2, and MMPs could serve as molecular targets for anti-metastatic therapies could be of paramount interest.

Previously, we demonstrated that TF-Ag mimicking and TF-Ag blocking inhibitors significantly reduce breast and prostate carcinoma metastasis *in vivo* by impeding the initial TF-Ag/Gal-3 mediated tumor cell adhesion to the endothelium [9, 19, 20]. Recently, yet another group used natural TF-Ag expressing glycopeptide TFD100 isolated from Atlantic cod to inhibit TF-Ag/Gal-3 mediated tumor/endothelial cell adhesion and ultimately PC-3 metastasis *in vivo* [31]. Inhibiting simultaneously additional subsequent steps of tumor metastasis mediated by endothelial integrins will increase dramatically our ability to control hematogenous spread of cancer. With these regards, it appears that endothelial $\alpha 3\beta 1$, Src, and MAP kinases could serve as valuable therapeutic targets. Further, the means for therapeutic targeting of the same signaling pathways in tumor cells are actively developed [32]. Identifying relevant molecular targets for anti-metastatic therapies located not in tumor cells, but in the target organ vasculature may present a new paradigm for controlling and preventing cancer metastasis.

MATERIALS AND METHODS

Cell lines and antibodies

Metastatic human prostate carcinoma PC-3 cells (ATCC, Rockville, MD) were maintained as monolayer cultures using RPMI-1640 media supplemented with 10% FBS in a 5% CO₂ humidified incubator. Human umbilical vein endothelial cells, HUVEC, (Life Technologies, Grand Island, NY) were cultured using Basal Medium 200 (Life Technologies) supplemented with low serum growth supplement containing FBS (2% v/v final concentration), hydrocortisone, human fibroblast growth factor, heparin, and human epidermal growth factor. The following antibodies were used in this study: anti-TF-Ag produced by JAA-F11 hybridoma [33]; anti-Gal-3 produced by TIB-166 hybridoma (ATCC); anti-integrin $\alpha 3$ (clone P1B5, EMD Millipore, Billerica, MA); Integrin Antibody Sampler Kit (#4749) including antibodies against integrins $\alpha 4$, $\alpha 5$, αV , $\beta 1$, $\beta 3$, $\beta 4$, and $\beta 5$; anti-phospho-Src (#2101); anti-Src (#2123); anti-phospho-p38 (#9211); anti-p38 (#9212); anti-phospho-MEK1/2 (#9121); anti-MEK1/2 (#9122); anti-phospho-Akt (#9271); anti-Akt (#9272) all from Cell Signaling, Danvers, MA; anti- β -Actin (ab8227) from Abcam, Cambridge, MA.

TF-Gold pull-down

Gold nanoparticles carrying multiple TF antigen epitopes (TF-Au) covalently attached to the particles through the polyethylene glycol (PEG) linker and control nanoparticles, on which PEG linkers are terminated with the OH groups (PEG-Au) and exhibiting excellent

solubility and stability were prepared as previously described [34]. Confluent endothelial cell monolayers grown for 6 days in collagen-coated T-150 flasks were treated for 60 min with TF-Au or PEG-Au nanoparticles (250 μ l of 4 mg/ml solution diluted in 20 ml of complete RPMI-1640 media containing 5% FBS), while endothelial cells treated with RPMI-1640 media containing 5% FBS only served as an additional control. Next, unbound nanoparticles were washed away by rinsing cultures twice with ice cold PBS; the cells were lysed using CellLytic-M mammalian cell lysis/extraction reagent (Sigma, Saint Louis, MO) supplemented with protease inhibitor cocktail (Sigma) and subjected to immunoprecipitation (IP) using JAA-F11 anti-TF-Ag antibody and protein A agarose. The immunoprecipitates were resolved on NuPAGE 4-12% gradient Bis-Tris gels (Invitrogen, Carlsbad, CA) and used either for Western blot analysis or LC-tandem mass spectrometry to identify precipitated proteins.

Western blot analysis

For TF-Gold pull-down analysis, TF-Au, PEG-Au, and untreated endothelial cell pull-down isolates were resolved on NuPAGE 4-12% gradient Bis-Tris gels (Invitrogen), and transferred to a nitrocellulose membrane (Invitrogen). The membranes were probed with primary antibodies against Gal-3, α 3, α 4, α 5, α V, β 1, β 3, β 4, and β 5 integrins, and Src kinase (see antibody list above) in conjunction with corresponding HRP-conjugated secondary antibodies and enhanced chemiluminescent (ECL) detection.

Endothelial cells from co-culture experiments (see below) were lysed using CellLytic M buffer with protein inhibitor cocktail (Sigma). Protein concentrations were determined using Protein Assay kit (Bio-Rad). Equal amounts of the protein from each sample (30 μ g) were resolved on a NuPAGE 4-12% gradient Bis-Tris gels (Invitrogen), and transferred to a nitrocellulose membranes (Invitrogen). The membranes were sequentially probed with primary antibodies directed against phosphorylated and total Src, p38, MEK1/2, and Akt (see antibody list above) in conjunction with corresponding HRP-conjugated secondary antibodies and enhanced chemiluminescent (ECL) detection. Anti- β -Actin antibody (Abcam, Cambridge, CA) was used to control loading. The experiments were performed at least three times for each setting with same results.

Mass spectrometry analysis

TF-Au, PEG-Au, and untreated endothelial cell pull-down isolates resolved on NuPAGE 4-12% gradient Bis-Tris gels were stained with modified colloidal Coomassie Blue G-250 [35] and 25 individual bands were cut by hand from each lane (Fig. 1, B). In gel digestion

using mass spectrometry grade trypsin gold (Promega, Madison, WI), peptide extraction, and LC-tandem mass spectrometry analyses on the Applied Biosystems-MDS-Sciex (Concorde, Ontario, Canada) 4000 Qtrap mass spectrometer were performed as previously described [36]. The tandem mass spectrometry data were processed for protein identifications using an in-house MASCOT search engine version 4.2 (Matrix Science, Boston, MA) using the Human NCBI protein database and one missed protease cleavage site. Variable modifications were allowed for oxidized methionines and a fixed modification for carbamidomethylated cysteines. Significant proteins hits were any protein(s) that had at least one individual peptide sequence score of >40 . The experiments were repeated twice with same results.

Modified parallel flow chamber experiments

A modification of the parallel flow chamber assay was developed to enable the analysis of changes in strength of adhesion between tumor and endothelial cells. Briefly, endothelial cells were grown until confluent in 35 mm tissue culture dishes. Next, 1 ml of a single tumor cell suspension (5×10^5 cell/ml) prepared immediately prior the experiment using nonenzymatic cell dissociation reagent (Sigma) was applied on top of the endothelial monolayer and allowed to interact with endothelial cells for various (5, 10, 15, 20, 30 min) time periods at 37°C in a CO₂ incubator (static adhesion phase). After that, unbound tumor cells were gently washed away, a parallel flow chamber deck (Glycotech, Rockville, MD) was mounted on top of the 35 mm dish on a stage of the inverted video microscope, and gradually increasing flow of a displacement media was initiated using a precision syringe pump KDS210 (KD Scientific, New Hope, PA) generating a wall shear force of a defined magnitude (parallel flow phase), while the process of cancer cell detachment and displacement was monitored and video recorded. Based on these experiments, temporal dynamics of tumor-endothelial cell adhesion stabilization were determined including: 1). The relationship between the time of tumor cell interaction with endothelium and a wall shear force necessary to displace 50% of tumor cells, and 2). The relationship between the wall shear force applied and a percentage of adherent tumor cells displaced. The experiments were performed at least three times for each setting and the data presented as means \pm STDEV.

Tumor cell/endothelial cell co-cultures

To investigate the changes in phosphorylation and expression of the proteins of interest in endothelial cells when they interact with tumor cells, an experimental system was developed enabling a fast and efficient separation of endothelial and cancer cells following

their interaction. To achieve this, we took advantage of the observation that endothelial cells, after growing on a collagen support for 6-7 days, become strongly attached to the collagen coated plastic and fairly resistant to the action of cell dissociation reagents, whereas tumor cells interacting with endothelium for a relatively short time (0-120 min) could be easily dislodged using non-enzymatic cell dissociation solution (Sigma). Thus, we performed series of experiments, in which 5×10^5 PC-3 cells stably transfected with GFP (for controlling the efficiency of tumor cell separation from endothelial cells) interacted for various periods of time (0, 30, 60, 120 min) with endothelial monolayers growing for 7 days in collagen-coated T150 flasks. After that tumor cells were quickly separated from endothelial monolayers using nonenzymatic cell dissociation reagent (Sigma), harvested, and frozen at -70°C for future analysis, while endothelial cells were lysed on plastic using cell lysis buffer with protein inhibitor cocktail (Sigma) and analyzed by Western blot as described above. The experiments were performed at least three times for each setting.

ACKNOWLEDGMENTS

This research was supported in parts by the Award Number 1I01BX000609 from the Biomedical Laboratory Research & Development Service of the VA Office of Research and Development (VVG) and the National Cancer Institute of the National Institutes of Health Award R01CA160461 (VVG). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health, or VA Office of Research and Development.

REFERENCES:

1. Labelle, M., Hynes, R.O. The initial hours of metastasis: the importance of cooperative host-tumor cell interactions during hematogenous dissemination. *Cancer Discov.* 2012; 2: 1091-1099.
2. Chambers, A.F., Groom, A.C., MacDonald, I.C. Dissemination and growth of cancer cells in metastatic sites. *Nat Rev Cancer.* 2002; 2: 563-572.
3. Glinsky, V.V. Intravascular cell-to-cell adhesive interactions and bone metastasis. *Cancer Metastasis Rev.* 2006; 25: 531-540.
4. Bendas, G., Borsig, L. Cancer cell adhesion and metastasis: selectins, integrins, and the inhibitory potential of heparins. *Int J Cell Biol.* 2012; Article ID 676731. doi: 10.1155/2012/676731.
5. Strell, C., Niggemann, B., Voss, M.J., Powe, D.G., Zänker, K.S., Entschladen, F. Norepinephrine promotes the $\beta 1$ -integrin-mediated adhesion of MDA-MB-231 cells to vascular endothelium by the induction of a $\text{GRO}\alpha$ release. *Mol Cancer Res.* 2012; 10: 197-207.
6. Senapati, S., Chaturvedi, P., Chaney, W.G., Chakraborty, S., Gnanapragassam, V.S., Sasson, A.R., Batra, S.K. Novel Interaction of MUC4 and galectin: potential pathobiological implications for metastasis in lethal pancreatic cancer. *Clin Cancer Res.* 2011; 17: 267-274.
7. Gassmann, P., Kang, M.L., Mees, S.T., Haier, J. In vivo tumor cell adhesion in the pulmonary microvasculature is exclusively mediated by tumor cell-endothelial cell interaction. *BMC Cancer.* 2010; 10: 177. doi: 10.1186/1471-2407-10-177.
8. Yu, L.G., Andrews, N., Zhao, Q., McKean, D., Williams, J.F., Connor, L.J., Gerasimenko, O.V., Hilkens, J., Hirabayashi, J., Kasai, K., Rhodes, J.M. Galectin-3 interaction with Thomsen-Friedenreich disaccharide on cancer-associated MUC1 causes increased cancer cell endothelial adhesion. *J Biol Chem.* 2007; 282: 773-781.
9. Glinskii, O.V., Huxley, V.H., Glinsky, G.V., Pienta, K.J., Raz, A., Glinsky, V.V. Mechanical Entrapment Is Insufficient and Intercellular Adhesion Is Essential for Metastatic Cell Arrest in Distant Organs. *Neoplasia.* 2005; 7: 522-527.
10. Dimitroff, C.J., Lechpammer, M., Long-Woodward, D., Kutok, J.L. Rolling of human bone-metastatic prostate tumor cells on human bone marrow endothelium under shear flow is mediated by E-selectin. *Cancer Res.* 2004; 64: 5261-5269.
11. Mine, S., Fujisaki, T., Kawahara, C., Tabata, T., Iida, T., Yasuda, M., Yoneda, T., Tanaka, Y. Hepatocyte growth factor enhances adhesion of breast cancer cells to endothelial cells in vitro through up-regulation of CD44. *Exp Cell Res.* 2003; 288: 189-197.
12. Draffin, J.E., McFarlane, S., Hill, A., Johnston, P.G., Waugh, D.J.J. CD44 potentiates the adherence of metastatic prostate and breast cancer cells to bone marrow endothelial cells. *Cancer Res.* 2004; 64: 5702-5711.
13. Havens, A.M., Jung, Y., Sun, Y.X., Taichman, R.S. The role of sialomucin CD164 (MGC-24v or endolyn) in prostate cancer metastasis. *BMC Cancer.* 2006; 6: 195. doi:10.1186/1471-2407-6-195.
14. Springer, G.F., Desai, P.R., Ghazizadeh, M., Tegtmeier, H. T/Tn pancarcinoma autoantigens: fundamental, diagnostic, and prognostic aspects. *Cancer Detect Prev.* 1995; 19: 173-182.
15. Almogren, A., Abdullah, J., Ghapure, K., Ferguson, K., Glinsky, V.V., Rittenhouse-Olson, K. Anti-Thomsen-Friedenreich-Ag (anti-TF-Ag) potential for cancer therapy. *Front Biosci (Schol Ed).* 2012; 4: 840-863.
16. Yu, L.G. The oncofetal Thomsen-Friedenreich carbohydrate antigen in cancer progression. *Glycoconj J.* 2007; 24: 411-420.
17. Glinsky, V.V., Glinsky, G.V., Rittenhouse-Olsen, K., Huflejt, M.E., Glinskii, O.V., Deutscher, S.L., Quinn, T.P. (2001) The Role of Thomsen-Friedenreich Antigen in Adhesion of Human Breast and Prostate Cancer Cells to

- the Endothelium. *Cancer Res.* 2001; 61: 4851-4857.
18. Khaldoyanidi, S.K., Glinsky, V.V., Sikora, L., Glinskii, A.B., Mossine, V.V., Quinn, T.P., Glinsky, G.V., Sriramarao, P. MDA-MB-435 human breast carcinoma cell homo- and heterotypic adhesion under flow conditions is mediated in part by Thomsen-Friedenreich antigen-galectin-3 interactions. *J Biol Chem.* 2003; 278: 4127-4134.
 19. Heimburg, J., Yan, J. Morey, S., Wild, L., Glinskii, O.V., Huxley, V.H., Klick, R., Roy, R., Glinsky, V.V., Rittenhouse-Olson, K. Inhibition of Spontaneous Breast Cancer Metastasis by Anti-Thomsen-Friedenreich Antigen Monoclonal Antibody JAA-F11. *Neoplasia.* 2006; 8: 939-948.
 20. Glinskii, O.V., Sud, S., Mossine, V.V., Mawhinney, T.P., Anthony, D.C., Glinsky, G.V., Pienta, K.J., Glinsky, V.V. Inhibition of Prostate Cancer Bone Metastasis by Synthetic TF Antigen Mimic/Galectin-3 Inhibitor Lactulose-L-Leucine. *Neoplasia.* 2012; 14: 65-73.
 21. Raz, A., Lotan, R. (1987) Endogenous galactoside-binding lectins: a new class of functional tumor cell surface molecules related to metastasis. *Cancer Metastasis Rev.* 1987; 6: 433-452.
 22. Inohara, H., Raz, A. (1994) Effects of natural complex carbohydrate (citrus pectin) on murine melanoma cell properties related to galectin-3 functions. *Glycoconj J.* 1994; 11: 527-532.
 23. Lotan, R., Raz, A. Endogenous lectins as mediators of tumor cell adhesion. *J Cell Biochem.* 1988; 37: 107-117.
 24. Nakahara, S., Oka, N., Raz, A. On the role of galectin-3 in cancer apoptosis. *Apoptosis.* 2005; 10: 267-275.
 25. Glinsky, V.V., Kiriakova, G., Glinskii, O.V., Mossine, V.V., Mawhinney, T.P., Turk, J.R., Glinskii, A.B., Huxley, V.H., Price, J.E., Glinsky, G.V. Synthetic Galectin-3 Inhibitor Increases Metastatic Cancer Cell Sensitivity to Taxol-Induced Apoptosis In Vitro and In Vivo. *Neoplasia.* 2009; 11: 901-909.
 26. Balan V, Wang Y, Nangia-Makker P, Kho D, Bajaj M, Smith D, Heilbrun L, Raz A, Heath E. Galectin-3: a possible complementary marker to the PSA blood test. *Oncotarget.* 2013; 4: 542-549.
 27. Glinskii, O.V., Turk, J.R., Pienta, K.J., Huxley, V.H., and Glinsky, V.V. Evidence of Porcine and Human Endothelium Activation by Cancer-Associated Carbohydrates Expressed on Glycoproteins and Tumor Cells. *J Physiol.* 2004; 554: 89-99.
 28. Fukushi, J., Makagiansar, I.T., Stallcup W.B: NG2 proteoglycan promotes endothelial cell motility and angiogenesis via engagement of galectin-3 and alpha3beta1 integrin. *Mol Biol Cell.* 2004; 15: 3580-3590.
 29. Takada, Y., Wayner, E.A., Carter, W.G., Hemler, M.E. (1988) Extracellular matrix receptors, ECMRII and ECMRI, for collagen and fibronectin correspond to VLA-2 and VLA-3 in the VLA family of heterodimers. *J Cell Biochem.* 1988; 37: 385-393.
 30. Munshi, H.G., Wu, Y.I., Mukhopadhyay, S., Ottaviano, A.J., Sassano, A., Koblinski, J.E., Platanias, L.C., Stack, M.S. Differential regulation of membrane type 1-matrix metalloproteinase activity by ERK 1/2- and p38 MAPK-modulated tissue inhibitor of metalloproteinases 2 expression controls transforming growth factor-beta1-induced pericellular collagenolysis. *J Biol Chem.* 2004; 279: 39042-39050.
 31. Guha, P., Kaptan, E., Bandyopadhyaya, G., Kaczanowska, S., Davila, E., Thompson, K., Martin, S.S., Kalvakolanu, D.V., Vasta, G.R., Ahmed, H. Cod glycopeptide with picomolar affinity to galectin-3 suppresses T-cell apoptosis and prostate cancer metastasis. *Proc Natl Acad Sci U S A.* 2013; 110: 5052-5057.
 32. McCubrey JA, Steelman LS, Chappell WH, Sun L, Davis NM, Abrams SL, Franklin RA, Cocco L, Evangelisti C, Chiarini F, Martelli AM, Libra M, Candido S, Ligresti G, Malaponte G, Mazzarino MC, Fagone P, Donia M, Nicoletti F, Polesel J, Talamini R, Bäsecke J, Mijatovic S, Maksimovic-Ivanic D, Michele M, Tafuri A, Dulińska-Litewka J, Laidler P, D'Assoro AB, Drobot L, Umezawa D, Montalto G, Cervello M, Demidenko ZN. Advances in targeting signal transduction pathways. *Oncotarget.* 2012; 3: 1505-1521.
 33. Rittenhouse-Diakun, K., Xia, Z., Pickhardt, D., Morey, S., Baek, M.-G., Roy, R. Development and characterization of monoclonal antibody to T-antigen: (Galβ1-3GalNAc-α-O). *Hybridoma.* 1998; 17: 165-173.
 34. Svarovsky, S.A., Szekely, Z., Barchi, J.J. Jr. Synthesis of gold nanoparticles bearing the Thomsen-Friedenreich disaccharide: A new multivalent presentation of an important tumor antigen. *Tetrahedr Asymm.* 2005; 16: 587-598.
 35. Candiano, G., Bruschi, M., Musante, L., Santucci, L., Ghiggeri, G.M., Carnemolla, B., Orecchia, P., Zardi, L., Righetti, P.G. Blue silver: a very sensitive colloidal Coomassie G-250 staining for proteome analysis. *Electrophoresis.* 25: 1327-1333.
 36. Li, F., Glinskii, O.V., Zhou, J., Wilson, L.S., Barnes, S., Anthony, D.C., Glinsky, V.V. Identification and Analysis of Signaling Networks Potentially Involved in Breast Carcinoma Metastasis to the Brain. *PLoS ONE.* 2011; 6(7): e21977.