Correction

Correction: Teneligliptin enhances the beneficial effects of GLP-1 in endothelial cells exposed to hyperglycemic conditions

Valeria De Nigris[1], Francesco Prattichizzo[2], Elettra Mancuso[1,3], Rosangela Spiga[1,3], Gemma Pujadas[1] and Antonio Ceriello[1,2,4]

[1] Institut d'Investigacions BiomeÌdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain

[2] Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) MultiMedica, Milan, Italy

[3] Department of Medical and Surgical Sciences, University Magna Grǽcia of Catanzaro, Catanzaro, Italy

[4] Centro de Investigacioìn Biomeìdica en Red de Diabetes y Enfermedades Metaboìlicas Asociadas (CIBERDEM), Madrid, Spain

***Published**: October 26, 2018*

Copyright: De Nigris et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been corrected: The correct affiliations are given below:

[1] Institut d'Investigacions BiomeÌdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain

[4] Centro de Investigacioìn Biomeìdica en Red de Diabetes y Enfermedades Metaboìlicas Asociadas (CIBERDEM), Madrid, Spain

Original article: Oncotarget. 2018; 9:8898-8910. https://doi.org/10.18632/oncotarget.22849