

Correction: PD-L1 expression heterogeneity in non-small cell lung cancer: evaluation of small biopsies reliability

Enrico Munari^{1,2}, Giuseppe Zamboni¹, Marcella Marconi¹, Marco Sommaggio¹, Matteo Brunelli², Guido Martignoni^{2,3}, George J. Netto⁴, Francesca Moretta⁵, Maria Cristina Mingari⁶, Matteo Salgarello⁷, Alberto Terzi⁸, Vincenzo Picece⁹, Carlo Pomari¹⁰, Gianluigi Lunardi⁹, Alberto Cavazza¹¹, Giulio Rossi¹², Lorenzo Moretta¹³ and Giuseppe Bogina¹

¹ Department of Pathology, Sacro Cuore Don Calabria Hospital, Negrar, Italy

² Department of Pathology AOUI, University of Verona, Verona, Italy

³ Department of Pathology, Pederzoli Hospital, Peschiera del Garda, Italy

⁴ Department of Pathology, The University of Alabama at Birmingham, Birmingham, AL, USA

⁵ Department of Laboratory Medicine, Sacro Cuore Don Calabria Hospital, Negrar, Italy

⁶ Department of Experimental Medicine (DIMES), University of Genoa, Genoa, Italy

⁷ Department of Nuclear Medicine, Sacro Cuore Don Calabria Hospital, Negrar, Italy

⁸ Department of Thoracic Surgery, Sacro Cuore Don Calabria Hospital, Negrar, Italy

⁹ Department of Oncology, Sacro Cuore Don Calabria Hospital, Negrar, Italy

¹⁰ Department of Pulmonology, Sacro Cuore Don Calabria Hospital, Negrar, Italy

¹¹ Department of Pathology, Arcispedale S. Maria Nuova/IRCCS, Reggio Emilia, Italy

¹² Department of Pathology, Azienda USL Valle d'Aosta, Aosta, Italy

¹³ Immunology Research Area, IRCCS Bambino Gesù Pediatric Hospital, Rome, Italy

Published: December 11, 2018

Copyright: Munari et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been corrected: In the Materials and Methods section, regarding the Tissue Microarray Construction, the diameter of the TMA core is incorrect. The proper value should be 1 mm, not 0.6 mm, as shown below:

Tissue microarray construction

For every case, all H&E stained slides were reviewed for diagnosis confirmation; one block was then selected for tissue microarray (TMA) construction. It has been demonstrated that TMAs containing at least 3 cores per case yield satisfactory agreement compared with whole section in lung cancer [14]. Therefore, for each block, 5 cores with a diameter of 1 mm were obtained randomly from the diverse areas of the tumor. Overall, 11 TMAs were built.

Original article: Oncotarget. 2017; 8:90123-90131. <https://doi.org/10.18632/oncotarget.21485>