Correction

Correction: Prediction of survival prognosis of non-small cell lung cancer by APE1 through regulation of epithelial-mesenchymal transition

Xi Wei^{1,*}, Qing Li^{2,*}, Ying Li^{3,*}, Wei Duan², Chongbiao Huang⁴, Xiangqian Zheng⁷, Lei Sun⁵, Jingtao Luo⁶, Dong Wang², Sheng Zhang¹, Xiaojie Xin¹ and Ming Gao⁷

Published: January 13, 2022

Copyright: © 2022 Wei et al. This is an open access article distributed under the terms of the <u>Creative Commons Attribution License</u> (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been corrected: In Figure 4D, the 3rd row, 3rd panel image is an accidental duplicate of the 1st row, 2nd panel image. In addition, the figure legend for 4D has been updated to read, 'D. A549 cells were indicated by immunofluorescence after APE1 siRNA and inhibitor AT101 treatment. (magnification: 1000×)'. The corrected Figure 4D, produced using the original data, is shown below. The authors declare that these corrections do not change the results or conclusions of this paper.

Original article: Oncotarget. 2016; 7:28523-28539. https://doi.org/10.18632/oncotarget.8660

¹Department of Diagnostic and Therapeutic Ultrasonography, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center of Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin, China

²Cancer Center, Daping Hospital and Research Institute of Surgery, Third Military Medical University, Chongqing, China

³The Third Department of Breast Cancer, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center of Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin, China

⁴Department of Senior Ward, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center of Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin, China

⁵Department of Biochemistry and Molecular Biology, Tianjin Medical University Cancer Institute and Hospital, Tianjin, China

⁶The Department of Otorhinolaryngology and Maxillofacial Oncology, Tianjin Medical University Cancer Institute & Hospital, Key Laboratory of Cancer Prevention and Therapy, Tianjin Cancer Institute, National Clinical Research Center of Cancer, Tianjin, China

⁷Department of Thyroid and Cervical Tumor, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center of Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin, China

^{*}These authors contributed equally to this work




Figure 4: (D) A549 cells were indicated by immunofluorescence after APE1 siRNA and inhibitor AT101 treatment. (magnification: 1000×).