

Correction

Correction: Berberine induces autophagy in glioblastoma by targeting the AMPK/mTOR/ULK1-pathway

Jiwei Wang¹, Qichao Qi¹, Zichao Feng¹, Xin Zhang¹, Bin Huang¹, Anjing Chen¹, Lars Prestegarden^{2,3}, Xingang Li¹ and Jian Wang^{1,2}

¹Department of Neurosurgery, Qilu Hospital of Shandong University and Brain Science Research Institute, Shandong University, Jinan, 250012, P.R. China

²Department of Biomedicine, University of Bergen, 5009 Bergen, Norway

³Department of Dermatology, University of Bergen, 5021 Bergen, Norway

Published:

Copyright: © 2021 Wang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/3.0/) (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been corrected: Due to errors during figure creation, the images of BBR 100 μM (0 h) in both U251 and U87 in Figure 3A contain overlapping areas (row 1, panels 1 and 2 and row 1, panels 3 and 4). The corrected Figure 3A, obtained using the original data, is shown below. The authors declare that these corrections do not change the results or conclusions of this paper.

Original article: Oncotarget. 2016; 7:66944–66958. <https://doi.org/10.18632/oncotarget.11396>




Figure 3: BBR impairs migration and invasion in GBM cells. (A) The migratory ability and statistical results of the migratory rate of U251 and U87 cells were evaluated by a wound healing assay treated with different concentrations of BBR for 24 h.