

Addendum

Addendum: Transfected SARS-CoV-2 spike DNA for mammalian cell expression inhibits p53 activation of p21(WAF1), TRAIL Death Receptor DR5 and MDM2 proteins in cancer cells and increases cancer cell viability after chemotherapy exposure

Shengliang Zhang^{1,2,3,4} and Wafik S. El-Deiry^{1,2,3,4,5}

¹Laboratory of Translational Oncology and Experimental Cancer Therapeutics, Warren Alpert Medical School, Brown University, Providence, RI 02912, USA

²Department of Pathology and Laboratory Medicine, Warren Alpert Medical School, Brown University, Providence, RI 02912, USA

³Joint Program in Cancer Biology, Lifespan Health System and Brown University, Providence, RI 02912, USA

⁴Legorreta Cancer Center at Brown University, Providence, RI 02912, USA

⁵Hematology/Oncology Division, Department of Medicine, Lifespan Health System and Brown University, Providence, RI 02912, USA

Published:

Copyright: © 2024 Zhang and El-Deiry. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#) (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has an addendum: This article received a publication fee waiver.

Original article: Oncotarget. 2024; 15:275–284. <https://doi.org/10.18632/oncotarget.28582>